



AN EFFICIENT TECHNIQUE IN FINDING THE EXACT SOLUTIONS FOR CAUCHY PROBLEMS

¹. DEPARTMENT OF ARCHITECTURAL ENGINEERING, SHOMAL UNIVERSITY, AMOL, IRAN

². DEPARTMENT OF ARCHITECTURAL ENGINEERING, SHAHID BEHESHTI UNIVERSITY, TEHRAN, IRAN

ABSTRACT: Nonlinear phenomena play a crucial role in applied mathematics and engineering. In this paper, the Reconstruction of Variational Iteration Method (RVIM) is used to solve the Cauchy problem. The new applied algorithm is a powerful and efficient technique in finding solutions for the linear and nonlinear equations using only few terms. The method used gives rapidly convergent successive approximations. As stated before, we aim to obtain analytical solutions to problems. We also aim to confirm that the reconstruction of variational iteration method is powerful, efficient, and promising in handling scientific and engineering problems. The Reconstruction of Variational Iteration Method (RVIM) technique is independent of any small parameters at all. Besides, it provides us with a simple way to ensure the convergence of solution series, so that we can always get accurate enough approximations. Some examples are given to elucidate the solution procedure and reliability of the obtained results.

KEYWORDS: Reconstruction of Variational Iteration Method, Cauchy problem, Inviscid Burgers' equation, Transport equation

INTRODUCTION

Nonlinear phenomena play a crucial role in applied mathematics and engineering. The theory of nonlinear problems has recently undergone many studies. Various approximation methods have been used for complex equations. In recent years, several such techniques have drawn special attention, such as Hirota's bilinear method [1], the homogeneous balance method [2-3], inverse scattering method [4], Adomian's decomposition method ADM [5,6], the variational iteration method [7,8] the \square -expansion method [9], as well as homotopy analysis method (HAM). After that, many types of nonlinear problems were solved by the HAM by others [10].

One of the newest approximation methods is Reconstruction of variational iteration method [11-15]. This new applied algorithm is a powerful and efficient technique in finding the approximate solutions for the linear and nonlinear equations. RVIM as a method that based on Laplace transform has high rapid convergence and reduces the size of calculations using only few terms, so many linear and nonlinear equations can be solved by this method. The method used gives rapidly convergent successive approximations. As stated before, we aim to obtain analytical solutions to problems. We also aim to confirm that the reconstruction of variational iteration method is powerful, efficient, and promising in handling scientific and engineering problems.

The RVIM technique is independent of any small parameters at all. Besides, it provides us with a simple way to ensure the convergence of solution series, so that we can always get accurate enough approximations. The RVIM technique has been successfully applied to many nonlinear problems in science and engineering. All of these verify the great potential and validity of the RVIM technique for strongly nonlinear problems in science and engineering.

In this paper, we use the method to discuss the first-order partial differential equation in the form [16]:

$$u_t(x,t) + a(x,t)u_x(x,t) = \psi(x), \quad x \in R, t > 0 \quad (1)$$

$$u(x,0) = \varphi(x), \quad x \in R \quad (2)$$

When $a(x,t) = a$ is a constant and $\psi(x) = 0$ Eq. (1) is a linear equation called the transport equation which can describe many interesting phenomena such as the spread of AIDS, the moving of wind.

When $a(x,t) = u(x,t)$ the equation is called the inviscid Burgers' equation arising in one-dimensional stream of particles or fluid having zero viscosity.

DESCRIPTION OF THE NEW METHOD

To clarify the basic ideas of our proposed method in [19], we consider the following differential equation same as VIM based on Lagrange multiplier [20]:

$$Lu(x_1, \dots, x_k) + Nu(x_1, \dots, x_k) = f(x_1, \dots, x_k) \quad (3)$$

By suppose that

$$Lu(x_1, \dots, x_k) = \sum_{i=0}^k L_{xi}u(x_i) \quad (4)$$

where L is a linear operator, N a nonlinear operator and f an inhomogeneous term.

We can rewrite equation (3) down a correction functional as follows:

$$L_{xj}u(x_j) = \underbrace{f(x_1, \dots, x_k) - Nu(x_1, \dots, x_k) - \sum_{\substack{i=0 \\ i \neq j}}^k L_{xi}u(x_i)}_{h((x_1, \dots, x_k), u(x_1, \dots, x_k))} \quad (5)$$

therefore

$$L_{xj}u(x_j) = h((x_1, \dots, x_k), u(x_1, \dots, x_k)) \quad (6)$$

With artificial initial conditions being zero regarding the independent variable x_j .

By taking Laplace transform of both sides of the equation (6) in the usual way and using the artificial initial conditions, we obtain the result as follows

$$P(s)U(x_1, \dots, x_{i-1}, s, x_{i+1}, x_k) = H((x_1, \dots, x_{i-1}, s, x_{i+1}, x_k), u) \quad (7)$$

where $P(s)$ is a polynomial with the degree of the highest derivative in equation (7), (the same as the highest order of the linear operator $L x_j$). The following relations are possible;

$$\ell[h] = H \quad (8-a)$$

$$B(s) = \frac{1}{P(s)} \quad (8-b)$$

$$\ell[b(x_i)] = B(s) \quad (8-c)$$

Which that in equation (8-a) the function $H((x_1, \dots, x_{i-1}, s, x_{i+1}, x_k), u)$ and $h((x_1, \dots, x_{i-1}, x_i, x_{i+1}, x_k), u)$ have been abbreviated as respectively.

Hence, rewrite the equation (7) as;

$$U(x_1, \dots, x_{i-1}, s, x_{i+1}, x_k) = H((x_1, \dots, x_{i-1}, s, x_{i+1}, x_k), u)B(s) \quad (9)$$

Now, by applying the inverse Laplace Transform on both sides of equation (9) and by using the (8-a) - (8-c), we have;

$$u(x_i, \dots, x_{i-1}, x_i, x_{i+1}, x_k) = \int_0^{x_i} h(x_1, \dots, x_{i-1}, \tau, x_{i+1}, x_k, u) b(x_i - \tau) d\tau \quad (10)$$

Now, we must impose the actual initial conditions to obtain the solution of the equation (3). Thus, we have the following iteration formulation:

$$u_{n+1}(x_1, \dots, x_{i-1}, x_i, x_{i+1}, x_k) = u_0(x_1, \dots, x_{i-1}, x_i, x_{i+1}, x_k) + \int_0^{x_i} h((x_1, \dots, x_{i-1}, \tau, x_{i+1}, x_k), u) b(x_i - \tau) d\tau \quad (11)$$

where u_0 is initial solution with or without unknown parameters.

Assuming u_0 is the solution of Lu with initial/boundary conditions of the main problem, In case of no unknown parameters, should satisfy initial/ boundary conditions. When some unknown parameters are involved in u_0 , the unknown parameters can be identified by initial/boundary conditions after few iterations, this technology is very effective in dealing with boundary problems. It is worth mentioning that, in fact, the Lagrange multiplier in the He's variational iteration method is $\lambda(\tau) = b(x_i - \tau)$ as shown in [19].

The initial values are usually used for selecting the zeroth approximation. With u_0 determined, then several approximations u_n $n > 0$, follow immediately. Consequently, the exact solution may be obtained by using

$$u(x_1, \dots, x_{i-1}, x_i, x_{i+1}, x_k) = \lim_{n \rightarrow \infty} u_n(x_1, \dots, x_{i-1}, x_i, x_{i+1}, x_k). \quad (12)$$

In what follows, we will apply the RVIM method to homogeneous/non-homogeneous parabolic partial differential equations to illustrate the strength of the method and to establish exact solutions for these problems.

APPLICATION

Since our focus is on the ideas and basic principles, we shall consider only the simplest possible equations to clearly illustrate the solution procedure. In particular, we will focus on pure Cauchy problems. These problems are initial value problems.

Example1. Consider the transport equation [16-18]:

$$u_t(x,t) + au_x(x,t) = 0, \quad x \in R, \quad t > 0 \quad (13)$$

Subject to the initial conditions:

$$u(x,0) = x^2, \quad x \in R \quad (14)$$

By selection of auxiliary linear operator, so the Eq. (13) is rewritten as

$$\ell\{u(x,t)\} = u_t = -\overbrace{(au_x)}^{h(x,\varepsilon,u)} \quad (15)$$

Now Laplace transform is implemented with respect to independent variable x on both sides of eq. (15) and by using the new artificial initial condition (which all of them are zero) we have

$$s \cup(x,t) = \ell\{h(x,t,u)\} \quad (16)$$

$$\cup(x,t) = \frac{\ell\{h(x,t,u)\}}{s} \quad (17)$$

and whereas Laplace inverse transform of $1/s$ is as follows

$$\ell^{-1}\left[\frac{1}{s}\right] = 1 \quad (18)$$

Therefore by using the Laplace inverse transform and convolution theorem it is concluded that

$$u(x,t) = \int_0^t h(x,\varepsilon,u) d\varepsilon \quad (19)$$

Therefore, by using the Eq. (10), (11) one can obtain the following RVIM's iteration formula in the t -direction

$$u_{n+1}(x,t) = u_0(x,t) - \int_0^t au_{nx} d\varepsilon \quad (20)$$

Now we start with an arbitrary initial approximation $u_0(x,t) = x^2$, that satisfies the initial condition and by using the RVIM iteration formula (20), we have the following successive approximation

$$u_1(x,t) = x^2 - 2atx + a^2t^2 \quad (21)$$

$$u_2(x,t) = x^2 - 2atx + a^2t^2 \quad (22)$$

$$u_3(x,t) = x^2 - 2atx + a^2t^2 \quad (23)$$

Whereas, the RVIM method admits the use of

$$u(x,t) = \lim_{n \rightarrow \infty} u_n(x,t),$$

which gives the exact solution

$$u(x,t) = x^2 - 2atx + a^2t^2 \quad (24)$$

Example 2. Consider the nonlinear Cauchy problem [17]

$$u_t(x,t) + xu_x(x,t) = 0, \quad x \in R, \quad t > 0 \quad (25)$$

Subject to the initial conditions:

$$u(x,0) = x^2, \quad x \in R \quad (26)$$

Such as previous examples, for implementation of the RVIM technique, first of all we need to choose the auxiliary linear operator as

$$\ell\{u(x,t)\} = u_t = -\overbrace{(xu_x)}^{h(x,\varepsilon,u)} \quad (27)$$

Accordingly, after taking Laplace Transform by using the artificial initial condition as in [19], on both side of Eq. (25), the following RVIM iteration formula in the t -direction can be obtained

$$u_{n+1}(x,t) = u_0(x,t) - \int_0^t xu_{nx} d\varepsilon \quad (28)$$

By the RVIM'S recurrent formula in Eq. (28), the terms of the sequence $\{u_n\}$ are constructed as follows, so that we choose its initial approximate solution as $u_0(x,t) = x^2$.

$$u_1(x,t) = x^2 - 2tx^2 \quad (29)$$

$$u_2(x,t) = x^2 - 2tx^2 + 2t^2x^2 \quad (30)$$

$$u_2(x,t) = x^2 - 2tx^2 + 2t^2x^2 + \frac{4}{3}t^3x^2 \quad (31)$$

The next terms of $\{u_n\}$ can be determined in a similar way and we can construct the n^{th} approximation as

$$u(x,t) = x^2 e^{-2t} \quad (32)$$

The approximation obtained by RVIM procedure converges to the exact solution.

Example3. Now we solve the following non-homogeneous Cauchy problem [17]

$$u_t(x,t) + u_x(x,t) = x, \quad x \in R, t > 0 \quad (33)$$

$$u(x,0) = e^x, \quad x \in R$$

To apply RVIM to this equation with initial and boundary conditions, according to (5) and (6), we have

$$\ell\{u(x,t)\} = u_t = \overbrace{(-u_x + x)}^{h(x,\varepsilon,u)} \quad (34)$$

Therefore, as previous Examples the RVIM iterative formula can be expressed as:

$$u_{n+1}(x,t) = u_0(x,t) - \int_0^t x u_{nx} d\varepsilon \quad (35)$$

With the aid of the initial approximation $u_0(x,t) = e^x$ and using the RVIM iteration, we can obtain directly the rest of the other components as follows

$$u_1(x,t) = e^x - te^x + tx \quad (36)$$

$$u_2(x,t) = e^x - te^x + tx - \frac{t^2}{2} + e^x t^2 \quad (37)$$

$$u_3(x,t) = e^x - te^x + tx - \frac{t^2}{2} + e^x t^2 - e^x \frac{t^6}{6} \quad (38)$$

The RVIM admits the use of $u(x,t) = \lim_{n \rightarrow \infty} u_n(x,t)$, which gives the exact solution

$$u(x,t) = t(x - \frac{t}{2}) + e^{x-t} \quad (39)$$

Example4. Consider the inviscid Burgers' equation [17, 18]

$$u_t(x,t) + u(x,t)u_x(x,t) = x, \quad x \in R, t > 0 \quad (40)$$

$$u(x,0) = x, \quad x \in R$$

To implementation of the RVIM method to this differential equation with initial and boundary conditions, according to (5) and (6), the auxiliary linear operator is selected as

$$\ell\{u(x,t)\} = u_t = \overbrace{-(uu_x)}^{h(x,\varepsilon,u)} \quad (41)$$

Therefore, as previous Examples the RVIM iterative formula can be expressed as:

$$u_{n+1}(x,t) = u_0(x,t) - \int_0^t uu_{nx} d\varepsilon \quad (42)$$

We start with an initial approximation $u_0(x,t) = x$; by the iteration formula (42), we can obtain the first few components as follows

$$u_1(x,t) = x - tx \quad (43)$$

$$u_2(x,t) = x - tx + t^2 x - \frac{t^3 x}{3} \quad (44)$$

$$u_3(x,t) = x - tx + t^2 x - \frac{t^3 x}{3} + \frac{2t^4 x}{3} - \frac{2t^5 x}{3} + \frac{2t^6 x}{9} - \frac{t^7 x}{63} \quad (45)$$

and the rest of the components are obtained using the iteration formula (41) and the solution in closed form is given by. Recall that $u(x,t) = \lim_{n \rightarrow \infty} u_n(x,t)$,

Consequently, the exact solutions

$$u(x,t) = \frac{x}{1-t}. \quad (46)$$

CONCLUSIONS

There are two main goals that we aimed for this work. The first is employing the powerful Reconstruction of the variational iteration to investigate Cauchy problems and the second one is showing the power of this method and its significant features. The two goals are achieved.

It is obvious that the method gives rapidly convergent successive approximations without any restrictive assumptions or transformation that may change the physical behavior of the problem. Reconstruction of the variational iteration gives several successive approximations through using the RVIM's iteration relation. Moreover, the RVIM reduces the size of calculations by not requiring the tedious Adomian polynomials, and hence the iteration is direct and straightforward. The RVIM uses the initial values for selecting the zeroth approximation, and boundary conditions, when given for bounded domains, can be used for justification only.

A clear conclusion can be drawn from the results that Reconstruction of the variational iteration method provides us an efficient tool to obtaining exact solutions of partial differential equations

REFERENCES

- [1.] X.B. Hu, Y.T. Wu, Application of the Hirota bilinear formalism to a new integrable differential-difference equation, *Phys. Lett. A* 246 (1998) 523-529.
- [2.] E. Fan, Two new applications of the homogeneous balance method, *Phys. Lett. A* 256 (2000) 353-357.
- [3.] M. Wang, Y. Zhou, Z. Li, Applications of a homogeneous balance method to exact solution of nonlinear equations in mathematical physics, *Phys. Lett. A* 216 (1996) 67-75.
- [4.] V.O. Vakhnenko, E.J. Parkes, A.J. Morrison, A Bäcklund transformation and the inverse scattering transform method for the generalized Vakhnenko equation, *Chaos Solitons Fractals* 17 (2003) 683-692.
- [5.] G. Adomian, Nonlinear dissipative wave equations, *Appl. Math. Lett.* 11 (1998) 125-126.
- [6.] G. Adomian, R. Rach, Modified decomposition solution of linear and nonlinear boundary-value problems, *Nonlinear Anal.* 23 (1994) 615-619.
- [7.] M.T. Darvishi, F. Khani, A. A. Soliman, The numerical simulation for stiff systems of ordinary differential equations, *Comput. Math. Appl.* 54 (2007) 1055-1063.
- [8.] A. Nikkar, M. Mighani, Application of VIM for Solving Seventh-Order Differential Equations” *American Journal of Computational and Applied Mathematics*, 2(1) (2012) 37-40
- [9.] A.V. Karmishin, A.I. Zhukov, V.G. Kolosov, *Methods of Dynamics Calculation and Testing for Thin-walled Structures*, Mashinostroyenie, Moscow, 1990.
- [10.] Y. Khan, R. Taghipour, M. Fallahian, A. Nikkar, A new approach to modified regularized long wave equation , *Neural Computing & Applications*, (26 July 2012), pp.1-7, doi. 10.1007/s00521-012-1077-0
- [11.] A. Nikkar, “A new approach for solving Gas dynamics equation.” *Acta Technica Corviniensis-Bulletin of Engineering*, 2012, 4:113-116.
- [12.] A. Nikkar, J. Vahidi, M. Jafarnejad Ghomi, M. Mighani, Reconstruction of variation Iteration Method for Solving Fifth Order Caudrey-Dodd-Gibbon (CDG) Equation, *International journal of Science and Engineering Investigations*, 2012, 1(6): 38-41.
- [13.] A. Nikkar, Z. Mighani, S.M. Saghebian, S.B. Nojabaei and M. Daie, Development and validation of an analytical method to the solution of modelling the pollution of a system of lakes. *Res. J. Appl. Sci. Eng. Technol.*, 5(1) (2013) 296-302.
- [14.] S. Ghasempour, J. Vahidi, A. Nikkar, M. Mighani, Analytical approach to some highly nonlinear equations by means of the RVIM, *Res. J. Appl. Sci. Eng. Technol.*, 5(1) (2013) 339-345.
- [15.] A. A. Imani, D. D. Ganji, Houman B. Rokni, H. Latifizadeh, E. Hesameddini, M. Hadi Rafiee, Approximate traveling wave solution for shallow water wave equation, *Applied Mathematical Modelling*, 36(4) (2012) 1550-1557.
- [16.] F. Khani, M. Ahmadzadeh Raji, H. Hamedi Nejad, Analytical solutions and efficiency of the nonlinear fin problem with temperature-dependent thermal conductivity and heat transfer coefficient, *Commun. Nonlinear Sci. Numer. Simul.* 14 (2009) 3327-3338.
- [17.] A. Molabahrani, F. Khani, The homotopy analysis method to solve the Burgers-Huxley equation, *Nonlinear Anal. RWA* 10 (2) (2009) 589-600.
- [18.] M. Sajid, I. Ahmad, T. Hayat, M. Ayub, Unsteady flow and heat transfer of a second grade fluid over a stretching sheet, *Commun. Nonlinear Sci. Numer. Simul.* 14 (1) (2009) 96-108.
- [19.] F. Khani, M. Ahmadzadeh Raji, S. Hamedi-Nezhad, A series solution of the fin problem with a temperature-dependent thermal conductivity, *Commun. Nonlinear Sci. Numer. Simul.* 14 (2009) 3007_3017.
- [20.] F. Khani, A. Farmany, M. Ahmadzadeh Raji, A. Aziz, F. Samadi, Analytic solution for heat transfer of a third grade viscoelastic fluid in non-Darcy porous media with thermophysical effects, *Commun. Nonlinear Sci. Numer. Simul.* (2009) doi:10.1016/j.cnsns.2009.01.031.
- [21.] F. Khani, A. Aziz, Thermal analysis of a longitudinal trapezoidal fin with temperature dependent thermal conductivity and heat transfer coefficient, *Commun. Nonlinear Sci. Numer. Simul* (2009), doi:10.1016 .2009.04.028.

- [22.] R.C. McOwen, *Partial Differential Equations: Methods and Applications*, Prentice Hall, Inc., 1996.
- [23.] A. Tveito, R. Winther, *Introduction to Partial Differential Equations*, Springer-Verlag, Berlin, Heidelberg, 2005.
- [24.] N.H. Asmar, *Partial Differential Equations with Fourier Series and Boundary Value Problems*, Prentice Hall, Inc., 2005.
- [25.] E. Hesameddini, H. Latifizadeh, *Reconstruction of Variational Iteration Algorithms using the Laplace Transform*, *Int. J. Nonlinear Sci. Numer. Simul.* 10 (2009)1377-1382.
- [26.] A.M. Wazwaz, *The variational iteration method: A powerful scheme for handling Linear and nonlinear diffusion equations*, *Computers and Mathematics with Applications*, (2007) 933-939.



ANNALS of Faculty Engineering Hunedoara



- International Journal of Engineering

copyright © UNIVERSITY POLITEHNICA TIMISOARA,
FACULTY OF ENGINEERING HUNEDOARA,
5, REVOLUTIEI, 331128, HUNEDOARA, ROMANIA
<http://annals.fih.upt.ro>