

ALLELOPATHIC WEED MANAGEMENT EFFICACY IN SUSTAINABLE AGRICULTURAL SYSTEMS

¹ INMA – Bucuresti, ROMANIA

Abstract: Weed control represents one of the major challenges in modern agricultural systems due to its negative impact on crop yields and the increasing costs of production. Allelopathy is an ecological strategy that utilizes natural compounds produced by plants (allelochemicals) to inhibit the germination, growth, and development of weeds, thus offering a sustainable alternative to chemical herbicides. This article provides a synthesis of the mechanisms of allelopathy, examples of plants with allelopathic potential, and their application as tools for weed management, discussing both the advantages and limitations of integrating this approach into sustainable agricultural systems. Recent research highlights the potential of allelopathy under both controlled and field conditions, emphasizing the need for practical validation and the adaptation of strategies to local pedoclimatic conditions. However, the transition from controlled laboratory bioassays to complex field environment often reveals significant ecological and technical limitations that hinder its widespread adoption. The efficacy of these natural phytotoxins is governed by a precarious balance of soil dynamics, microbial degradation, and climatic fluctuations, often resulting in inconsistent weed control.

Keywords: allelopathy; weeds; sustainable management; ecological agriculture; allelochemicals

1. INTRODUCTION

Allelopathy refers to the direct or indirect effect exerted by plants on neighboring plants or on their associated microflora and microfauna through the production of allelochemicals that interfere with plant growth and development. The allelochemicals released by plants act as a defense system against microbial attacks, herbivore predation, or competition from other plants [1]. The study of allelopathy is a sub-discipline of chemical ecology that focuses on the effects of chemicals produced by plants or microorganisms on the growth and development of other plants in natural or agricultural systems [2]. These effects can be either positive or negative on the growth of surrounding plants.

The term “allelopathy” is derived from two distinct Greek words: allelon, meaning “mutual” or “each other,” and pathos, meaning “to suffer” or “feeling.” Although the term “allelopathie” was first used by the Austrian scientist Hans Molisch in 1937, chemical interactions between plants have been known for thousands of years. For instance, in 300 B.C., the Greek botanist Theophrastus mentioned the negative effects of chickpea on other plants, and later, Pliny the Elder, a Roman scholar of the 1st century A.D., noted the inhibitory effect of the walnut tree (*Juglans spp.*) on nearby crops.

Allelochemicals are released from different parts of plants through processes such as leaching from leaves or plant residues deposited on the soil, root exudation, volatilization from aerial organs, decomposition of plant residues, and other processes occurring in natural and agricultural systems [3]. Once released, allelochemicals can suppress the germination, growth, and establishment of surrounding plants or modify soil properties in the rhizosphere by influencing the microbial community [4,5]. Considering the important role of allelopathic substances in regulating plant communities, they can also be used as natural biodegradable herbicides, providing a sustainable alternative to synthetic herbicides [6-8].

Research on allelopathy has traditionally focused on assessing the phytotoxic activities of plant residues or crude plant extracts [9]. Recent studies in the field of allelopathy have made significant progress due to the rapid development of separation and structural elucidation techniques, which allow for the detection, isolation, and characterization of active compounds [10,11]. Allelochemicals are produced by plants as secondary metabolites or by microorganisms through decomposition processes.

Allelochemicals comprise a chemically diverse group of secondary metabolites, which can be classified into several categories based on their chemical structure and mode of action, as discussed in detail in the following sections.

Plant growth regulators, such as salicylic acid, gibberellic acid, and ethylene, are also considered allelochemicals. Allelochemicals vary in their mode of action, uptake, and effectiveness [12]. The mode of action of many identified allelochemicals is still not fully understood. Many allelochemicals exhibit mechanisms that are not utilized by synthetic herbicides, providing researchers with insights for the discovery of new modes of action [13]. Although the efficacy and specificity of many allelochemicals remain unknown or limited [14], they represent a viable alternative to synthetic herbicides. Numerous plant species have been identified worldwide for their allelopathic effects, and among these, several have been highlighted as having high potential for future research.

However, must be considered that changing the classical pesticides with allelopathic options may uncover profound ecological and technical barriers that impede large-scale implementation. The efficacy of these biogenic phytotoxins is perpetually modulated by a volatile intersection of soil edaphic properties, rapid microbial catabolism, and stochastic climatic events, which frequently culminate in erratic weed control performance. Beyond inconsistent suppression, the latent risk of residual phytotoxicity toward subsequent cash crops introduces a precarious variable into farm economics, potentially compromising germination rates and total biomass accumulation. The complexity is further compounded by the metabolic plasticity of both the donor and target species; for instance, the concentration of allelochemicals often fluctuates in response to localized environmental stressors, such as nutrient deficiencies or herbivory, making "dosage" nearly impossible to standardize in a field setting. Moreover, the rhizosphere acts as a complex biochemical reactor where adsorption to clay minerals or organic matter can sequester active molecules before they reach the target radicle. There is also the burgeoning concern of evolutionary adaptation, as persistent selection pressure from specific allelopathic cultivars may induce metabolic resistance within weed populations, paralleling the challenges faced with synthetic herbicides. Consequently, a comprehensive academic understanding of these multifaceted constraints is imperative. Only through the synthesis of metabolomics, soil science, and precision agronomy can researchers refine these natural interactions into reliable components of an integrated weed management (IWM) system that is both ecologically resilient and economically competitive in modern sustainable agriculture.

2. MATERIALS AND METHODS

This analysis is based on a systematic review of recent scientific literature concerning the use of allelopathy for weed control in agricultural systems. The selected studies include experimental research conducted in laboratories, greenhouses, and field conditions, as well as reviews documenting the effectiveness of allelopathic plants and their allelochemicals against various weed species.

MATERIALS

The materials analyzed for weed control through allelopathy include:

- Plant species with allelopathic potential: Rye (*Secale cereale*), Sorghum (*Sorghum bicolor*), Sunflower (*Helianthus annuus*), Indian Mustard (*Brassica juncea*), and Rice (*Oryza sativa*, allelopathic genotypes).
- Natural allelochemicals: phenols, terpenoids, flavonoids, organic acids, isothiocyanates, and sorgoleone.
- Plant-derived materials: plant residues used as mulch and plant extracts applied to the soil or on crop surfaces.
- Cover crops: planted as an intermediate layer to suppress weed germination and growth prior to the main crop.

METHODS

The methods for applying allelopathic materials for weed management include:

1. Cover crops - planting allelopathic species prior to the main crop to reduce weed germination and growth.
2. Plant mulch - using residues of allelopathic plants to gradually release allelochemicals into the soil.
3. Intercropping - simultaneously cultivating allelopathic plants with the main crop, combining allelopathic effects with direct competition for resources.
4. Biofumigation - incorporating allelopathic plants, such as *Brassica* spp., into the soil to release isothiocyanates and other inhibitory compounds.

5. Application of plant extracts - administering concentrated solutions from allelopathic plants directly into the soil or onto the crop surface to reduce weed germination and growth. The effectiveness of each method depends on the plant species used, the type of allelochemicals, crop density, mode of application, and pedoclimatic conditions, making it necessary to adapt strategies to the specific characteristics of each farm or region.

■ CLASSIFICATION OF ALLELOCHEMICALS

Allelochemicals are classified into 14 categories based on their chemical similarities. These categories include: water-soluble organic acids; straight-chain alcohols; aliphatic aldehydes and ketones; simple unsaturated lactones; long-chain fatty acids and polyacetylenes; benzoquinones, anthraquinones, and complex quinones; simple phenols, benzoic acid and its derivatives; cinnamic acid and its derivatives; coumarins; flavonoids; tannins; terpenoids and steroids; amino acids and peptides; alkaloids and cyanohydrins; sulfides and glucosinolates; as well as purines and nucleosides (Figure 1) [15].




Figure 1. Representative allelochemicals from the 14 categories classified based on their chemical similarities [15].

3. RESULTS AND DISCUSSION

■ MECHANISMS OF ALLELOPATHY

Allelopathy occurs through the release of allelochemicals into the soil, air, or water, affecting the physiological processes of weeds. These compounds can:

- Reduce the growth of roots and aerial parts;
- Disrupt essential metabolic processes.

Plants release these compounds through roots, decomposition of plant residues, or volatilization. The effectiveness of allelopathic interactions depends on environmental factors and soil type.

Allelopathic interactions involve the release of bioactive biochemical compounds into the surrounding environment, which interfere with the physiological and biochemical processes of weeds. The main pathways of allelochemical release and their effects on weed growth and development are schematically presented in Figure 2.

Figure 2 illustrates the mechanisms by which allelopathic plants release allelochemicals into the surrounding environment through root exudates, decomposition of plant residues, volatilization, and leaching from aerial organs. These compounds affect weed seed germination, root and shoot growth, nutrient uptake, and essential physiological processes, leading to reduced weed competitiveness against crop plants.




Figure 2. Main pathways of allelochemical release and their effects on weed control.

EXAMPLES OF PLANTS WITH ALLELOPATHIC POTENTIAL

- Recent research has identified various plant species with significant effects on weeds, including:
- Rye (*Secale cereale*) - suppresses the germination of annual weed species;
 - Sorghum (*Sorghum bicolor*) - produces the allelochemical sorgoleone, which has an inhibitory effect;
 - Cereal crops such as wheat, barley, and rice - exhibit allelopathic potential against problematic weeds, such as *Lolium rigidum* and *Portulaca oleracea*.

To highlight the main plant species with allelopathic potential investigated in the scientific literature and their use in weed management, Table 1 presents a comparative summary of allelopathic plants, the allelochemicals involved, their effects on weeds, and corresponding agronomic applications. These species can be integrated into sustainable agricultural systems through the use of cover crops, plant mulches, or biofumigation, contributing to reduced weed pressure and decreased reliance on chemical herbicides.

Table 1. Comparative summary of allelopathic plants, their allelochemicals, effects on weeds, and agronomic applications

Allelopathic Species	Main Allelochemicals	Effect on Weeds	Agronomic Application	Reference
Rye (<i>Secale cereale</i> L.)	Phenolic acids, benzoxazinoids	Inhibition of annual weed germination	Cover crop, plant mulch	[16]
Sorghum (<i>Sorghum bicolor</i> (L.) Moench)	Sorgoleone	Reduction of root growth	Mulch, plant extracts	[16,17]
Sunflower (<i>Helianthus annuus</i> L.)	Phenolic acids, flavonoids	Decreased weed vigor	Incorporation of plant residues into soil	[17]
Indian mustard (<i>Brassica juncea</i> (L.) Czern.)	Isothiocyanates	Inhibition of germination and growth	Biofumigation, green manure	[18]
Rice (<i>Oryza sativa</i> L., allelopathic genotypes)	Phenolics, organic acids	Reduction of weed density	Selective cultivation	[16,19]

The allelopathic effectiveness depends on pedoclimatic conditions, crop density, and the management of plant residues.

APPLICATION OF ALLELOPATHY IN AGRICULTURAL SYSTEMS

- Allelopathy can be integrated into weed management through:
- Plant mulches that gradually release allelochemicals into the soil;
- Intercropping with allelopathic species;
- Natural extracts applied as bioherbicides.

As summarized in Table 1, the use of allelopathic species as cover crops or sources of plant mulch can lead to a significant reduction in weed density, with the effect depending on the species, pedoclimatic conditions, and the applied cropping technology.

Studies have shown a significant reduction in weed density following the implementation of these practices; however, field validation is essential to calibrate optimal doses and strategies.

ECOLOGICAL AND TECHNICAL CONSTRAINTS OF ALLELOPATHIC WEED MANAGEMENT IN SUSTAINABLE AGROECOSYSTEMS

The fundamental challenge of implementing allelopathy in commercial agriculture is the inherent lack of species-specific selectivity, which stands in stark contrast to the targeted precision of modern synthetic herbicides. Many allelopathic cover crops, such as *Secale cereale* (cereal rye) or

Sorghum bicolor, release broad-spectrum inhibitors that may suppress small-seeded cash crops with the same efficacy as the intended weed targets, thereby risking significant stand reduction and uneven crop establishment. This issue is further exacerbated by the phenomenon of autotoxicity, or "soil sickness," wherein the accumulation of specific secondary metabolites inhibits the germination and vigor of the donor species' own subsequent generations. Such chemical feedback loops necessitate extended fallow periods or complex rotational design, which can disrupt the economic continuity of a production system and complicate long-term management strategies [19]. Once released, biogenic phytotoxins often bind to clay minerals or organic matter, sequestering them and making them unavailable to weed roots. This is worsened by soil microbes that rapidly break down these molecules or unpredictably transform them into different metabolites. Additionally, fluctuations in soil pH, temperature, and moisture accelerate the degradation of these natural compounds, leading to inconsistent performance in the field [20]. From a quantitative perspective, the efficacy of allelopathy is hindered by the rigorous requirement of maintaining a minimum inhibitory concentration within the soil solution at the exact phenological moment of weed germination. This concentration is frequently compromised by dilution effects, as excessive precipitation or irrigation can leach water-soluble allelochemicals beyond the active weed seed zone, allowing resilient or deep-rooted species to escape suppression entirely. Unlike synthetic formulations engineered for slow release or environmental stability, natural metabolites are often highly volatile or susceptible to photolysis, resulting in an exceptionally narrow window of bioactivity. This volatility makes it difficult for producers to ensure that the "natural dose" is sufficient to achieve commercially acceptable levels of control, often leading to patchy weed distribution and the necessity for supplemental mechanical intervention [21, 22]. The reliability of allelopathic systems is further undermined by significant genetic and varietal variability, as the biosynthesis of defensive metabolites is a highly plastic trait influenced by both the plant's genomic architecture and its response to localized environmental stressors. This metabolic cost often creates a physiological trade-off where high allelochemical production may divert carbon resources away from primary growth, potentially resulting in reduced crop yields or diminished biomass accumulation [23]. Furthermore, plants under stress from drought or nutrient deficiency may spike toxin production unpredictably, while those in optimal conditions may produce insufficient quantities for weed suppression, rendering the standardization of "allelopathic cultivars" a significant breeding challenge. Over time, these systems also face the risk of evolutionary adaptation; persistent selection pressure from specific biogenic toxins can induce the development of detoxification mechanisms within weed populations, paralleling the metabolic resistance pathways observed in response to synthetic herbicides and threatening the long-term sustainability of the biological approach [24-26].

4. CONCLUSIONS

Allelopathy cannot fully replace chemical herbicides at present, but it represents an essential component of integrated weed management strategies. It provides a promising tool for controlling weeds in an ecological and sustainable manner, reducing reliance on synthetic herbicides and contributing to the diversification of agronomic management strategies. The integration of allelopathic plants into cropping systems, along with the use of mulches and natural extracts, can enhance the efficiency and sustainability of weed management. Future research should focus on the practical validation of effects observed in the laboratory and on determining the optimal implementation methods across different regions and soil types.


In conclusion, while allelopathy represents a sophisticated biological mechanism for weed suppression, its integration into sustainable agriculture is currently hindered by the unpredictable nature of the soil-plant-microbe interface. The transition from the laboratory to the field exposes these natural phytotoxins to complex degradation pathways, fluctuating environmental conditions, and a fundamental lack of species selectivity that can jeopardize both crop health and economic stability. Achieving a reliable "biological dosage" remains a primary technical hurdle, as the volatile and transient nature of secondary metabolites often fails to provide the consistent pressure required for modern production standards. Furthermore, the potential for weeds to develop metabolic resistance and the inherent physiological costs to the donor plant suggest that allelopathy cannot be viewed as a standalone solution. Instead, its future utility lies in its refinement through advanced breeding and precision management, serving as one component of a diversified, multi-tactic integrated weed management strategy that balances ecological integrity with agricultural productivity.

Acknowledgement

This research was supported by Project PN 23 04 02 01, Contract no.: 9N/ 01.01.2023 SUSTAIN–DIGI –AGRI, Innovative biofertilizer production technology used to restore soil biodiversity and reduce the effects of drought on agricultural lands. This paper was supported by the project ADER cod.2.1.4/ 20.07.2023 „Cercetări privind perfecționarea managementului integrat de combaterea buruienilor la principalele culturi de câmp în contextul schimbărilor climatice și a restricțiilor Green Deal privind exploatarea durabilă a resurselor de mediu”.

Bibliography

- [1] Kong, C. H., Xuan, T. D., Khanh, T. D., Tran, H. D., & Trung, N. T. (2019). Allelochemicals and signaling chemicals in plants. *Molecules*, 24(15), 2737
- [2] Einhellig, F. A. (1995). Allelopathy: current status and future goals. *Allelopathy: Organism, Process, and Applications* Inderjit, K. M. M. D., Einhellig, F. A.. ACS Symposium Series (Washington, DC, USA: ACS Symposium Series 582; American Chemical Society) pp. 1–24.
- [3] Anaya, A. L. (1999). Allelopathy as a tool in the management of biotic resources in agroecosystems. *Crit. Rev. Plant Sci.* 18(6), 697–739
- [4] Weir, T. L., Park, S. W., Vivanco, J. M. (2004). Biochemical and physiological mechanisms mediated by allelochemicals. *Curr. Opin. Plant Biol.* 7(4), 472–479
- [5] Zhou, B., Kong, C. H., Li, Y. H., Wang, P., Xu, X. H. (2013). Crabgrass (*Digitaria sanguinalis*) allelochemicals that interfere with crop growth and the soil microbial community. *J. Agric. Food Chem.* 61 (22), 5310–5317.
- [6] Duke, S., Romagni, J., Dayan, F. (2000). Natural products as sources for new mechanisms of herbicidal action. *Crop Prot.* 19 (8-10), 583–589
- [7] Vyyyan, J. R. (2002). Allelochemicals as leads for new herbicides and agrochemicals. *Tetrahedron* 58 (9), 1631–1646
- [8] Khamare Y., Chen J., Marble S.C. (2022). Allelopathy and its application as a weed management tool: A review. *Front. Plant Sci., Sec. Functional Plant Ecology*, Volume 13.
- [9] Weston, L. A. (1996). Utilization of allelopathy for weed management in agroecosystems. *Agron. J.* 88 (6), 860–866
- [10] Mallik, A. U. (2000). Challenges and opportunities in allelopathy research: a brief overview. *J. Chem. Ecol.* 26 (9), 2007–2009
- [11] Scavo, A., Restuccia, A., Lombardo, S., Fontanazza, S., Abbate, C., Pandino, G., et al. (2020). Improving soil health, weed management and nitrogen dynamics by trifolium subterraneum cover cropping. *Agron. Sustain. Dev.* 40 (3), 1–12
- [12] Weston, L. A., Duke, S. O. (2003). Weed and crop allelopathy. *Crit. Rev. Plant Sci.* 22 (3-4), 367–389.
- [13] Duke, S., Dayan, F., Rimando, A., Schrader, K., Aliotta, G., Oliva, A., et al. (2002). Invited paper: Chemicals from nature for weed management. *Weed Sci.* 50 (2), 138–151
- [14] Bhadoria, P. (2010). Allelopathy: A natural way towards weed management. *Am. J. Exp. Agric.* 1 (1), 7–20.
- [15] Cheng, F., Cheng, Z. (2015). Research progress on the use of plant allelopathy in agriculture and the physiological and ecological mechanisms of allelopathy. *Front. In Plant Sci.* 6.
- [16] Sahrir M.A.S., Yusoff N., Azizan K.A. (2023). Allelopathy activity under laboratory, greenhouse and field conditions: A review. *AIMS Agriculture and Food*, 8(1): 78–104
- [17] Murthy I.S., Jolly G.E., John A.P. (2025). Allelopathy in Weed Management: A Comprehensive Review. *International Journal of Plant & Soil Science*, 37(5):96–104
- [18] González-García, E., Sánchez-Moreiras A.M., Vieites-Álvarez Y. (2025) Allelopathic potential and chemical profile of wheat, rice and barley against herbicide-resistant weeds *Portulaca oleracea* L. and *Lolium rigidum* Gaud. *BMC Plant Biology*.
- [19] Parte V., Chouhan M., Nayak K. (2025) Allelopathic approaches for weed management. *International Journal of Advanced Biochemistry Research*, Vol. 9, Special Issue 2.
- [20] Scavo, A., & Mauromicale, G. (2021). Crop allelopathy for sustainable weed management in agroecosystems: Knowing the present with a view to the future. *Agronomy*, 11(11), 2104.
- [21] Ain, Q., Mushtaq, W., Shadab, M., & Siddiqui, M. B. (2023). Allelopathy: An alternative tool for sustainable agriculture. *Physiology and Molecular Biology of Plants*, 29(4), 495–511
- [22] Scavo, A.; Restuccia, A.; Mauromicale, G. Allelopathy: General principles and basic aspects for agroecosystem control. In *Sustainable Agriculture Reviews*; Gaba, S., Smith, B., Lichtfouse, E., Eds.; Springer: Cham, Switzerland, 2018; Volume 28, pp. 47–101
- [23] Khanh, T.D.; Chung, M.I.; Xuan, T.D.; Tawata, S. (2005). The exploitation of crop allelopathy in sustainable agricultural production. *J. Agron. Crop. Sci.* 191, 172–184
- [24] Bhowmik, P.C.; Inderjit. (2003) Challenges and opportunities in implementing allelopathy for natural weed management. *Crop. Prot.* 22, 661–671
- [25] Rial, C.; Tomé, S.; Varela, R.M.; Molinillo, J.M.G.; Macías, F.A. (2020). Phytochemical study of safflower roots (*Carthamus tinctorius*) on the induction of parasitic plant germination and weed control. *J. Chem. Ecol.*, 46, 871–880
- [26] Tlili, N.; Nasri, N.; Saadaoui, E.; Khaldi, A.; Triki, S. (2009). Carotenoid and tocopherol composition of leaves, buds and flowers of *Capparis spinosa* grown wild in Tunisia. *J. Agric. Food Chem.*, 57, 5381–5385.



ISSN 1584 – 2665 (printed version); ISSN 2601 – 2332 (online); ISSN-L 1584 – 2665
copyright © University POLITEHNICA Timisoara, Faculty of Engineering Hunedoara,
5, Revolutiei, 331128, Hunedoara, ROMANIA
<http://annals.fih.upt.ro>